

Cuts here,
This blue section does not
print

Acceleron® B-300 SAT applied? If "Yes" then Acceleron B-300 SAT, a biological, has been applied to this seed.	Yes	No
Acceleron B-300 SAT Batch #:		
Acceleron B-300 SAT not claimed effective after:		
Store seed in cool, dry conditions. To ensure efficacy of Acceleron B-300 SAT after the season of purchase, store seed at ≤ 68°F/20°C. Contains microorganisms. Microorganisms have the potential to elicit allergic reactions. For sensitized individuals, use appropriate PPE to reduce exposure to treated seed.		

CODE	SEED-APPLIED TECHNOLOGY
A	Metalaxyl
F	Prothioconazole
H1	Clothianidin at 0.25 mg ai/seed rate
H2	Clothianidin at 0.50 mg ai/seed rate
H3	Clothianidin at 1.25 mg ai/seed rate
I	Trifloxystrobin
L	Fluoxastrobin
N	Tioxazafen
Q	LCO SP104
V	<i>Bacillus firmus</i> I-1582
W	Ipconazole

WARNING: TREATED SEED. DO NOT USE TREATED SEED FOR FOOD, FEED OR OIL PURPOSES.

This seed has been treated at the manufacturer's recommended rate with the products identified by code on this seed tag, and decoded in the table in this section.

General Information: Handle treated seeds in ventilated areas. Avoid breathing or making contact with dust. If contaminated, remove clothing and laundry before reuse. Wash hands and face before eating, drinking, or smoking.

Personal Protective Equipment: When opening this bag, or loading, pouring, or handling the treated seed or seed pieces, wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made out of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride (PVC) ≥14 mils, or viton ≥14 mils.

First Aid: Eyes: Hold eye open and rinse gently with water for 15 minutes. If wearing contact lenses, remove after first 5 minutes and continue rinsing. **Skin or clothing:** Remove contaminated clothing. Rinse skin immediately with plenty of water. Use soap if available. **Swallowed:** Call poison center or doctor immediately for treatment advice. Sip water if able to swallow. Do not induce vomiting unless told to do so by the poison center or doctor. Do not give anything by mouth to an unconscious person. **Inhaled:** Move to fresh air. If not breathing, call 911 or ambulance, and then give artificial respiration. Call poison center or doctor for further treatment advice. **FOR EMERGENCY MEDICAL TREATMENT INFORMATION, CALL COLLECT DAY OR NIGHT: (314) 694-4000.**

Hazards to Humans and Animals: Harmful to humans if swallowed, inhaled or absorbed through skin. Treated seed is toxic to birds and mammals. Exposed treated seed is hazardous to wildlife, birds and mammals that come in contact with treated seed. Cover or collect treated seeds that are spilled during loading and planting, and in areas such as row ends.

Storage and Disposal: Store treated seed away from food and feed, and do not allow access by children, pets, or livestock. Do not reuse or refill empty seed bags. Surplus treated seed or empty treated seed containers should be stored or disposed according to local requirements. Dispose of all excess treated seed by burying away from bodies of water. Do not contaminate bodies of water when disposing of excess treated seed or wash waters of planting equipment. Dispose of seed and seed packaging or containers in accordance with local requirements. Do not use empty seed bags for any other purpose. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Excess treated seed may be used for biofuel (including ethanol) production only if: (1) By-products are not used for livestock feed, and (2) No measurable residues of pesticides remain in ethanol by-products that are used for agronomic practice.

PRODUCT SPECIFIC ADDITIONAL INFORMATION:

CLOTHIANIDIN + BACILLUS FIRMUS I-1582-TREATED SEED: This seed has been treated with PONCHO®/VOTIVO®, which contains 40.3% clothianidin and 8.1% *Bacillus firmus* I-1582. This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust, which can drift to blooming crops or weeds. To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's recommendations. Use planting equipment that will plant treated seed into the soil to a minimum depth of 0.5 inch. PONCHO/VOTIVO is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert. PONCHO/VOTIVO is not to be used in areas where rice/crawfish aquaculture practices are in place or in or near fish farms, shrimp, prawn or crab ponds or nursery operations. Regardless of type of application (seed treatment or foliar), do not apply more than 0.2 lbs (Soybean), 0.1 lbs (corn), 0.2 lbs (cotton), 0.14 lbs (sugarbeet), or 0.02 lbs (sorghum) of active ingredient clothianidin per acre per year. **Immediate Plant-back:** Cereal grains, field corn, popcorn, sweet corn, rapeseed, canola, soybeans, root and tuber vegetables **30-Day Plant-back:** Grasses, non-grass animal feeds, dry beans, dry peas; **8-Month Plant-back:** Sugar cane, **12-Month Plant-back:** Any crops without an earlier plant-back interval.

LCO SP104 TREATED SEED: There are no restrictions on crop rotation.

METALAXYL TREATED SEED: Acceleron DX-309 Fungicide Seed Treatments is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium*, and *Phytophthora* spp. Do not use with other seed treatment products unless previous experience assures compatibility.

PROTHIOCONAZOLE-TREATED SEED: Treated areas may be replanted with any crop specified on this label as soon as practical after last application. For crops not listed on this label, do not plant back within 30 days of application. After the seeds have been planted, do not enter or allow worker entry into the treated areas during the restricted-entry interval (REI) of 12 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with treated seeds in soil or with the planting media. To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's recommendations. Treated seeds must be planted at a minimum depth of 0.5 inches.

TIOXAZAFEN-TREATED SEED: Do not use at a rate that will result in more than 0.28 lb tioxazafen per acre per year. In case of early season crop failure, seeds (treated or untreated) may be immediately replanted as long as the maximum yearly application rate per acre is not exceeded. Treated seed must be planted into the soil at a depth greater than 1 inch. Do not plant seeds by broadcasting to the soil surface. **Plant-back Restrictions:** Corn, cotton, and soybeans may be replanted immediately. All other crops may be planted after 30 days.

TRIFLOXYSTROBIN-TREATED SEED: Areas planted with treated seed may be replanted immediately with any crop for which a tolerance exists for trifloxystrobin. Do not plant any crop without trifloxystrobin tolerances until 30 days after planting trifloxystrobin treated seed. After the seeds have been planted, do not enter or allow worker entry into the treated areas during the restricted-entry interval (REI) of 12 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with treated seeds in soil or with the planting media. To reduce seed dust which can drift onto blooming crops or weeds, ensure that planting equipment is functioning properly in accordance with manufacturer's recommendations.

Acceleron® and NemaStrike™ are trademarks of Monsanto Technology LLC. Poncho® and VOTIVO® are registered trademarks of Bayer.

©2010-2018 MONSANTO COMPANY All Rights Reserved

13089S2-291

