



Treating Site ID Number	Treatment Batch Number
Date Seed Treatment Applied	Quantity of Seed Treated:
	Units
Customer	

Acceleron® B-200 SAT applied?	Yes	No
If “Yes” then Acceleron B-200 SAT, a biological, has been applied to this seed.		
Acceleron B-200 SAT Batch #:		

Cuts here,  
This blue section does not  
print





<b>Treating Site ID Number</b>	<b>Treatment Batch Number</b>
<b>Date Seed Treatment Applied</b>	<b>Quantity of Seed Treated:</b>
	Units
<b>Customer</b>	

<b>Acceleron® B-200 SAT applied?</b> If “Yes” then Acceleron B-200 SAT, a biological, has been applied to this seed.	<b>Yes</b>	<b>No</b>
<b>Acceleron B-200 SAT Batch #:</b>		

## **WARNING:** TREATED SEED. DO NOT USE TREATED SEED FOR FOOD, FEED OR OIL PURPOSES

This seed has been treated at the manufacturer's recommended rate with Fluxapyroxad, Imidacloprid, Metalaxyl, Pyraclostrobin, and Tioxazafen. If Acceleron B-200 SAT is checked “yes”, a biological has also been applied to this seed.

**General Information:** Handle treated seeds in ventilated areas. Avoid breathing or making contact with dust. If contaminated, remove clothing and launder before reuse. Wash hands and face before eating, drinking, or smoking.

**Personal Protective Equipment:** When opening this bag or loading, pouring, or handling the treated seed or seed pieces, wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or viton ≥ 14 mils.

**Hazards to Humans and Animals:** Harmful to humans if swallowed, inhaled, or absorbed through skin. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds that spilled during loading, planting, and in areas such as row ends.

**FIRST AID: Eyes:** Hold eye open and rinse gently with water for 15 minutes. If wearing contact lenses, remove after first 5 minutes and continue rinsing. **Skin or clothing:** Remove contaminated clothing. Rinse skin immediately with plenty of water. Use soap if available. **Swallowed:** Call poison center or doctor immediately for treatment advice. Sip water if able to swallow. Do not induce vomiting unless told to do so by the poison center or doctor. Do not give anything by mouth to an unconscious person. **Inhaled:** Move to fresh air. If not breathing, call 911 or ambulance, and then give artificial respiration. Call poison center or doctor for further treatment advice. **For emergency medical treatment information, Call Collect Day or Night: (314) 694-4000.**

**Storage and Disposal:** Store treated seed away from food and feed, and do not allow access by children, pets, or livestock. Do not reuse or refill empty seed bags. Surplus treated seed or empty treated seed containers should be stored or disposed according to local requirements. Do not contaminate bodies of water when disposing of excess treated seed or wash waters of planting equipment. Dispose of them in accordance with local requirements. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Excess treated seed may be used for ethanol production only if: (1) by-products are not used for livestock feed, and (2) no measurable residues of pesticides remain in ethanol by-products that are used in agronomic practice.

**Restrictions:** Treated seed must be planted into the soil at a depth equal to or greater than 1 inch. Plant treated seed into the soil to the recommended minimum depth to minimize exposure. Metalaxyl is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium*, and *Phytophthora* spp. Do not use with other seed treatment products unless previous experience assures compatibility. After the seeds, have been planted, do not enter or allow worker entry into the treated areas during the restricted-entry interval (REI) of 12 hours. Exception: Once the seeds are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with treated seeds in soil or with the planting media. Plant-back Restrictions: Corn, cotton, and soybeans may be replanted immediately. After 30 days the following may be replanted: barley, brassica vegetables, bulb vegetables, cucurbits, dried shelled peas and beans, edible podded legume vegetables, flax, fruiting vegetables, leafy vegetables, mint, oat, rapeseed (includes canola and crambe), root vegetables, rye, sorghum, succulent shelled peas and beans, sugar beet, sunflower, tuberous and corn vegetables (including potato), wheat and triticale. Do not plant any other crop in treated area for 12 months are treated seeds are planted. In case of early season crop failure, seeds (treated or untreated) may be immediately replanted as long as the maximum yearly application rate per acre is not exceeded. Do not use at a rate that will result in more than 0.28 lb tioxazafen per acre per year.

Acceleron® and NemaStrike™ are trademarks of Monsanto Technology LLC. NemaStrike™  
Monsanto Company, 800 N. Lindbergh Blvd. - St. Louis, MO 63167  
©2018 MONSANTO COMPANY All Rights Reserved

13116S4-30





**ACCELERON<sup>®</sup>**  
SEED APPLIED SOLUTIONS

**ELITE**

WITH **NEMASTRIKE<sup>™</sup>**  
TECHNOLOGY