

JumpStart[®] wettable powder

phosphate-solubilizing inoculant

In-furrow application peanut:

- Add the contents of one 57 g (2.0 oz) bottle, one 80 g (2.8 oz) bottle, one 285 g (10.0 oz), or one 400 g (14.0 oz) container to the appropriate amount of water and apply at planting.
- Once JumpStart is mixed in water, apply to seed in-furrow.
 1. **Clean the liquid system with a cleaning solution**, JumpStart is a biological product, which may be sensitive to chemical residues
 2. **Convert the sprayer to in-furrow application.** Remove spray tips and replace with appropriate flow regulator "orifice discs". Proper orifice disc number may be determined from spraying manual. The objective is to obtain a solid stream of liquid into the furrow. Place spraying assembly behind the seed planting assembly so that the solid stream of product will flow into the furrow and on the seed. JumpStart should be on the seed and in the furrow before it closes, not to the side nor above or below the seed.
 3. **Fill clean spray tank half full with clean, non-chlorinated water.** Pull the JumpStart package from the carton, remove the cap and empty into the inoculant bag and shake well. Add remainder of clean, non-chlorinated water to spray tank to the appropriate solution level. Use the diluted product within 24 hours. Do not allow the diluted tank mix to exceed 100°F.
 4. **Application rate/unit coverage JumpStart is applied in-furrow at rates on table 3 below.** The quantity of JumpStart applied per acre varies depending upon the row width. The measurement chart provides the recommended rate of JumpStart per acre and also indicates the number of acres treated with each row width. The application of the diluted liquid should be approximately 3 to 20 gallons/acre based on equipment used and soil moisture. Calibrate sprayer to deliver correct amount.

Table 3: In-furrow applications rates for peanut

ROW WIDTH	APPLICATION RATE	57 g (2 OZ) BOTTLE TREATS	80 g (2.8 oz) BOTTLE TREATS	285 g (10 OZ) BOTTLE TREATS	400 g (14 OZ) BOTTLE TREATS
76.2 cm (30")					
Single row	11.6 g/ha (0.166 oz/ac)	3.4 ha (8.3 acres)	4.71 ha (11.6 acres)	16.8 ha (41.5 acres)	23.5 ha (58.1 acres)
Twin row	23.3 g/ha (0.332 oz/ac)	1.7 ha (4.2 acres)	2.36 ha (5.8 acres)	8.5 ha (21.0 acres)	11.9 ha (29.4 acres)
91.4 cm (36")					
Single row	14.0 g/ha (0.2 oz/ac)	4.04 ha (10 acres)	5.67 ha (14 acres)	20.2 ha (50 acres)	28.4 ha (70 acres)
Twin row	28.0 g/ha (0.4 oz/ac)	2.02 ha (5 acres)	2.83 ha (7 acres)	10.1 ha (25 acres)	14.2 ha (35 acres)

Where to use for maximum benefit

For soils low to medium in available phosphate:

Use JumpStart with the lower recommended P fertilizer rate from soil test results. If you do not soil test, use with your normal P fertilizer rate.

For soils high to very high in available phosphate:

Replace the starter in-row P fertilizer application [16.8-22.4 kg P₂O₅/ha (15-20 lb P₂O₅/acre)] with JumpStart.

Benefits of using JumpStart may be limited on:

- extremely sandy soils (greater than 85% sand).
- extremely high organic matter soils (greater than 14% organic matter).
- fields that have been heavily manured over the last several years.

Applicators and Other Handlers

The active ingredient in JumpStart phosphate-solubilizing inoculant is *Penicillium bilaiae* spores. Eye or skin contact, inhalation, or ingestion should be avoided. Wear standard protective clothing and equipment including gloves, safety glasses. Ensure adequate ventilation. Where exposure through inhalation may occur, use respiratory protective equipment. This product contains microorganisms that may have the potential to provoke sensitizing reactions; use appropriate personal protective equipment (PPE) to reduce exposure. Sensitized individuals should wear a NIOSH-approved respirator. In case of contact with skin or eyes, immediately flush exposed areas with plenty of water. Get medical attention if irritation occurs.

Storage and Disposal

The active ingredient in JumpStart is a living organism and requires specific storage conditions to ensure viability and product performance. To maintain product viability:

- Store JumpStart containers and seed to which JumpStart has been applied in a cool, unheated facility <20°C (68°F) away from sunlight and direct heat sources.
- Minimize temperature fluctuations.
- Avoid freeze/thaw cycles.
- Use entire contents of container once opened.
- Use before the expiry date. The expiry date is valid only for unopened containers stored under the conditions listed above.
- Do not mix with unapproved products.
- For a list of compatible seed treatments, contact Monsanto Company at 1-877-775-8787.

Unused JumpStart and unplanted seed to which JumpStart has been applied should be disposed of in accordance with applicable Federal, state/provincial, and local requirements.

Limited Warranty

The Seller warrants that this product contains a minimum number of *Penicillium bilaiae* colony forming units as specified on this label. The Seller makes no other warranty expressed or implied as to product viability or performance since product storage, use and growing conditions are beyond the Seller's control. Seller's guarantee is limited to the terms set out on the label and subject thereto. Buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition. If this product does not perform as warranted above, and to the extent consistent with applicable law, customer's sole remedy for breach of warranty shall be replacement of the product or refund of the purchase price paid, at the option of Monsanto Company.

EXCEPT AS PROVIDED ELSEWHERE IN WRITING CONTAINING AN EXPRESSED REFERENCE TO THIS WARRANTY AND LIMITATION OF DAMAGES, SELLER MAKES NO OTHER EXPRESSED OR IMPLIED WARRANTY OR GUARANTEE TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, INCLUDING ANY OTHER EXPRESSED OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO.

JumpStart[®] and Monsanto BioAg and Design[™] are trademarks of Monsanto Technology LLC. Used under license. ©2016 Monsanto Company. All rights reserved. Registration Number 2015029A Fertilizers Act. **Product of Canada.** 114029R4-39



JumpStart[®] wettable powder

phosphate-solubilizing inoculant

For use on: alfalfa, canola, chickpea, corn, dry bean, lentil, mustard, pea, peanut, sorghum, soybean, sugar beet, sunflower, sweetclover, and all wheat

ATTENTION:
This Specimen label is provided for general information only.

- This biological product may not yet be available or approved for sale or use in your area.
- It is your responsibility to follow all Federal, state/provincial and local laws and regulations regarding the use of biologicals.
- Before using any biological, be sure the intended use is approved in your state/province or locality.
- Your state/province or municipality may require additional precautions and instructions for use of this product that are not included here.
- Monsanto does not guarantee the completeness or accuracy of this Specimen label. The information found in this label may differ from the information found on the product label.
- You should not base any use of a similar product on the precautions, instructions for use or other information you find here.
- Always follow the precautions and instructions for use on the label of the biological you are using.

MINIMUM GUARANTEED ANALYSIS

ACTIVE: 7.2 x 10⁸ cfu/g *Penicillium bilaiae*

INERT: wettable powder

NON PLANT FOOD INGREDIENT

Not a fertilizer substitute

Read this entire label before using the product. Use only according to label directions.

How it Works

The active ingredient in JumpStart[®] phosphate-solubilizing inoculant is the soil fungus *Penicillium bilaiae*, which grows on plant roots and makes less-available residual soil phosphate available for crop use.

Directions for Use

On-seed treatment (all crops except peanut):

Inoculate seed on-farm by adding the contents of one 57 g (2.0 oz) bottle, one 80 g (2.8 oz) bottle, one 285 g (10.0 oz), or one 400 g (14.0 oz) container to the appropriate amount of water (refer to Table 1) prior to or during seeding.

JumpStart can be applied utilizing commercial on seed application equipment. Applicators used previously for pesticides should be triple rinsed before using for JumpStart application.

To divide the contents of a bottle of JumpStart, first suspend entire contents in 500 ml (0.5 US quarts) water, mix thoroughly and divide the mixture into smaller quantities as desired. JumpStart in solution must be applied to the seed within 24 hours.

JumpStart can be used with all nitrogen-fixing rhizobial inoculants. To mix JumpStart with a liquid inoculant, first suspend the entire contents of the JumpStart wettable powder bottle in a small amount of the liquid inoculant for easier mixing. Pour the mixture into the remaining liquid Rhizobia inoculant and mix to suspend. No additional water is required.

Continuous flow application

Wheat, pea, lentil, chickpea, or dry bean

- Apply the JumpStart suspension to seed using a flow-regulated applicator when transferring seed from the bin to the truck, or from the truck to the tank or seed cart.
- Fill the applicator tank with the required volume of cool, clean water (refer to Table 1).
- While mixing continuously, slowly add the contents of the JumpStart container to the water in the applicator tank.
- Mix well and agitate continuously to avoid settling.
- Determine the conveyor or auger flow rate in bu/min, and set the applicator to achieve the required flow rate (refer to Table 2). For best mixing, run the auger or conveyor at full speed and less than full capacity.

Batch application

Corn, soybean, sunflower, sugar beet, sorghum, canola, mustard, alfalfa, or sweet clover

- Apply the JumpStart suspension to seed using a batch treater or cement mixer.
- Fill the applicator tank with the required volume of cool, clean water (refer to Table 1).
- Add the contents of one 57 g (2.0 oz) bottle, one 80 g (2.8 oz) bottle, one 285 g (10.0 oz), or one 400 g (14.0 oz) JumpStart container to the water in the applicator tank.
- Mix well and agitate continuously to avoid settling.
- Put seed in the batch treater or cement mixer. Do not overload treater with seed. There must be enough space for mixing.
- Slowly apply the proper volume of JumpStart suspension to the seed as it tumbles (approximately 2 minutes). Example: if you are treating 1 bag/25 kg (55 lb) of alfalfa seed from a 57 g (2.0 oz) container, use 1/3 of the suspension.
- Re-bag the inoculated seed.
- Repeat with the remaining bags of seed. Note re-shake or continue agitating the JumpStart suspension in the applicator until fully suspended before inoculating the remaining bags of seed.



Manufactured For and Guaranteed By: Monsanto Company 800 N. Lindbergh Blvd. St. Louis, Missouri 63167 USA

For Medical, Transportation, Spill or Other Emergencies Call Collect 24 Hours a Day 1-314-694-4000

For Product Use Information Call: 1-877-775-8787

JumpStart[®] wettable powder

phosphate-solubilizing inoculant

JumpStart[®] wettable powder

phosphate-solubilizing inoculant

Table 1: On-seed application rates and bare seed planting windows

57 g (2.0 oz) container			
Crop	Seed inoculated by one 57 g (2.0 oz) container	Approximate water volume	Planting window (bare seed)
Wheat	1,100 kg (2,400 lb, 40 bu)	7.0 litres (7.6 US quarts)	30 days
Pea	1,900 kg (4,200 lb, 70 bu)	6.0 litres (6.3 US quarts)	30 days
Chickpea	1,500 kg (3,300 lb, 55 bu)	4.0 litres (4.2 US quarts)	30 days
Dry Bean ¹	1,100 kg (2,400 lb, 40 bu)	3.5 litres (3.7 US quarts)	30 days
Lentil	1,100 kg (2,400 lb, 40 bu)	3.5 litres (3.7 US quarts)	30 days
Soybean	1,135 kg (2,500 lb, 42 bu)	3.5 litres (3.7 US quarts)	30 days
Sunflower ²	100 kg (230 lbs)	2.0 litres (2.1 US quarts)	30 days
Sorghum ³	136 kg (300 lbs)	1.8 litres (1.9 US quarts)	60 days
Alfalfa/Sweet Clover	68 kg (150 lbs)	1.5 litres (1.6 US quarts)	7 days
Canola/Mustard	63 kg (140 lbs)	1.5 litres (1.6 US quarts)	60 days
Sugar beet ³	5 units	1.0 litres (1.0 US quarts)	7 days
Corn	800,000 kernels (10 bags ⁴)	2.8 litres (3.0 US quarts)	60 days

80 g (2.8 oz) container			
Crop	Seed inoculated by one 80 g (2.8 oz) container	Approximate water volume	Planting window (bare seed)
Wheat	1,630 kg (3,600 lb, 60 bu)	10.0 litres (11.0 US quarts)	30 days
Pea	2,720 kg (6,000 lb, 100 bu)	8.0 litres (8.5 US quarts)	30 days
Chickpea	2,180 kg (4,800 lb, 80 bu)	6.0 litres (6.4 US quarts)	30 days
Dry Bean ¹	1,630 kg (3,600 lb, 60 bu)	5.0 litres (5.3 US quarts)	30 days
Lentil	1,630 kg (3,600 lb, 60 bu)	5.0 litres (5.3 US quarts)	30 days
Soybean	1,630 kg (3,600 lb, 60 bu)	5.0 litres (5.3 US quarts)	30 days
Sunflower ²	140 kg (320 lbs)	2.8 litres (3.0 US quarts)	30 days
Sorghum ³	190.4 kg (420 lbs)	2.5 litres (2.7 US quarts)	60 days
Alfalfa/Sweet Clover	100 kg (220 lbs)	2.0 litres (2.2 US quarts)	7 days
Canola/Mustard	90 kg (198 lbs)	2.0 litres (2.2 US quarts)	60 days
Sugar beet ³	7 units	1.4 litres (1.5 US quarts)	7 days
Corn	1,120,000 kernels (14 bags ⁴)	3.92 litres (4.14 US quarts)	60 days

285 g (10.0 oz) container			
Crop	Seed inoculated by one 285 g (10.0 oz) container	Approximate water volume	Planting window (bare seed)
Wheat	5,500 kg (12,000 lb, 200 bu)	35.0 litres (38.0 US quarts)	30 days
Pea	9,500 kg (21,000 lb, 350 bu)	30.0 litres (31.0 US quarts)	30 days
Chickpea	7,500 kg (16,500 lb, 275 bu)	20.0 litres (23.0 US quarts)	30 days
Dry Bean ¹	5,500 kg (12,000 lb, 200 bu)	17.5 litres (20.0 US quarts)	30 days
Lentil	5,500 kg (12,000 lb, 200 bu)	17.5 litres (20.0 US quarts)	30 days
Soybean	5,675 kg (12,500 lb, 208 bu)	17.5 litres (20.0 US quarts)	30 days
Sunflower ²	500 kg (1,150 lb)	10.0 litres (11.0 US quarts)	30 days
Sorghum ³	680 kg (1,500 lb)	9.0 litres (9.5 US quarts)	60 days
Alfalfa/Sweet Clover	340 kg (750 lb)	7.5 litres (8.0 US quarts)	7 days
Canola/Mustard	315 kg (700 lb)	7.5 litres (8.0 US quarts)	60 days
Sugar beet ³	25 units	4.7 litres (5.0 US quarts)	7 days
Corn	4,000,000 kernels (50 bags ⁴)	14.0 litres (15.0 US quarts)	60 days

400 g (14 oz) container			
Crop	Seed inoculated by one 400 g (14.0 oz) container	Approximate water volume	Planting window (bare seed)
Wheat	8,150 kg (18,000 lb, 300 bu)	50.0 litres (53.0 US quarts)	30 days
Pea	13,600 kg (30,000 lb, 500 bu)	40.0 litres (42.0 US quarts)	30 days
Chickpea	10,900 kg (24,000 lb, 80 bu)	30.0 litres (32.0 US quarts)	30 days
Dry Bean ¹	8,150 kg (18,000 lb, 300 bu)	25.0 litres (27.0 US quarts)	30 days
Lentil	8,150 kg (18,000 lb, 300 bu)	25.0 litres (27.0 US quarts)	30 days
Soybean	8,150 kg (18,000 lb, 300 bu)	25.0 litres (27.0 US quarts)	30 days
Sunflower ²	700 kg (1540 lbs)	14.0 litres (15.0 US quarts)	30 days
Sorghum ³	952 kg (2094 lbs)	12.6 litres (13.5 US quarts)	60 days
Alfalfa/Sweet Clover	500 kg (1100 lbs)	10.0 litres (11.0 US quarts)	7 days
Canola/Mustard	455 kg (1000 lbs)	10.0 litres (11.0 US quarts)	60 days
Sugar beet ³	35 units	7 litres (7.5 US quarts)	7 days
Corn	5,600,000 kernels (70 bags ⁴)	19.6 litres (20.71 US quarts)	60 days

¹ Use with pinto, great northern, black, navy, kidney, red, and pink bean.
² When treating confectionary sunflowers, increase water rates to ensure proper coverage.
³ The application rate is currently under review. The rate provided is based on our experience with other crops.
⁴ 80,000 kernels per bag.

- Once JumpStart is mixed in water, apply to seed within 24 hours.
- The planting window is the maximum time allowed between JumpStart application and seeding.
- Application with other seed treatments is possible but may reduce the planting window. For up-to-date pesticide compatibility information and bare seed planting windows, please contact Monsanto Company at 1-877-775-8787.

Table 2. Applicator flow rate calibration (based on one 57 g (2.0 oz) container or 1/5 285 g (10 oz) container)

Auger flow rate				Applicator rate ¹									
				Wheat		Pea		Lentil and Dry Bean		Chickpea		Soybean	
bu/hr	bu/min	lb/hr	lb/min	minutes/ container ²	quarts/ min	minutes/ container ²	quarts/ min	minutes/ container ²	quarts/ min	minutes/ container ²	quarts/ min	minutes/ container ²	quarts/ min
240	4	14,400	240	10	0.75	17.5	0.34	10	0.38	13.8	0.33	10.3	0.37
360	6	21,600	360	6.7	1.13	11.7	0.51	6.7	0.56	9.2	0.49	6.8	0.55
480	8	28,800	480	5	1.5	8.8	0.69	5	0.75	6.9	0.65	5.1	0.73
600	10	36,000	600	4	1.88	7	0.86	4	0.94	5.5	0.82	4.1	0.92
720	12	43,200	720	3.3	2.25	5.8	1.03	3.3	1.13	4.6	0.98	3.4	1.1
840	14	50,400	840	2.9	2.63	5	1.2	2.9	1.31	3.9	1.15	2.9	1.28
960	16	57,600	960	2.5	3	4.4	1.37	2.5	1.5	3.4	1.31	2.6	1.46
1080	18	64,800	1080	2.2	3.38	3.9	1.54	2.2	1.69	3.1	1.47	2.3	1.65
1200	20	72,000	1200	2	3.75	3.5	1.71	2	1.88	2.8	1.64	2.1	1.83

Auger flow rate				Applicator rate ¹									
				Wheat		Pea		Lentil and Dry Bean		Chickpea		Soybean	
bu/hr	bu/min	kg/hr	lb/min	minutes/ container ²	L/min	minutes/ container ²	L/min	minutes/ container ²	L/min	minutes/ container ²	L/min	minutes/ container ¹	L/min
240	4	6,532	109	10	0.71	17.5	0.32	10	0.36	13.8	0.31	10.3	0.35
360	6	9,798	163	6.7	1.07	11.7	0.48	6.7	0.53	9.2	0.46	6.8	0.52
480	8	13,063	218	5	1.42	8.8	0.65	5	0.71	6.9	0.62	5.1	0.69
600	10	16,329	272	4	1.78	7	0.81	4	0.89	5.5	0.78	4.1	0.87
720	12	19,595	327	3.3	2.13	5.8	0.97	3.3	1.07	4.6	0.93	3.4	1.04
840	14	22,861	381	2.9	2.49	5	1.14	2.9	1.24	3.9	1.09	2.9	1.21
960	16	26,127	435	2.5	2.84	4.4	1.30	2.5	1.42	3.4	1.24	2.6	1.38
1080	18	29,393	490	2.2	3.20	3.9	1.46	2.2	1.60	3.1	1.39	2.3	1.56
1200	20	32,659	544	2	3.55	3.5	1.62	2	1.78	2.8	1.55	2.1	1.73

¹ Applicator rates are calculated assuming one 57 g (2.0 oz) container or 1/5 285 g (10 oz) container of JumpStart is suspended in 7 L (7.6 US quarts) of water to inoculate 40 bu of wheat, 6.0 L (6.3 US quarts) of water to inoculate 70 bu of pea, 4.0 L (4.2 US quarts) of water to inoculate 55 bu of chickpea, 3.5 L (3.7 US quarts) of water to inoculate 41 bu of soybean, or 3.5 L (3.7 US quarts) of water to inoculate 40 bu of lentil or dry bean.

² Minutes to apply one 57 g (2.0 oz) container or 1/5 285 g (10 oz) container of JumpStart. (See instructions for dividing container contents in Directions for Use).